

CAMS Suicide Status Form (SSF-IV-R) Initial Session

Patient: _____ Clinician: _____ Date: _____ Time: _____

Section A (Patient):

Rate and fill out each item according to how you feel right now.

Rank Then rank in order of importance 1 to 5 (1=most important to 5=least important).

_____	1) RATE PSYCHOLOGICAL PAIN (<i>hurt, anguish, or misery in your mind; <u>not</u> stress, <u>not</u> physical pain</i>): Low pain: 1 2 3 4 5 :High pain What I find most painful is: _____
_____	2) RATE STRESS (<i>your general feeling of being pressured or overwhelmed</i>): Low stress: 1 2 3 4 5 :High stress What I find most stressful is: _____
_____	3) RATE AGITATION (<i>emotional urgency; feeling that you need to take action; <u>not</u> irritation; <u>not</u> annoyance</i>): Low agitation: 1 2 3 4 5 :High agitation I most need to take action when: _____
_____	4) RATE HOPELESSNESS (<i>your expectation that things will not get better no matter what you do</i>): Low hopelessness: 1 2 3 4 5 :High hopelessness I am most hopeless about: _____
_____	5) RATE SELF-HATE (<i>your general feeling of disliking yourself; having no self-esteem; having no self-respect</i>): Low self-hate: 1 2 3 4 5 :High self-hate What I hate most about myself is: _____
N/A	6) RATE OVERALL RISK OF SUICIDE: Extremely low risk: 1 2 3 4 5 :Extremely high risk (will <u>not</u> kill self) (will kill self)

1) How much is being suicidal related to thoughts and feelings about yourself? Not at all: 1 2 3 4 5 : completely

2) How much is being suicidal related to thoughts and feelings about others? Not at all: 1 2 3 4 5 : completely

Please list your reasons for wanting to live and your reasons for wanting to die. Then rank in order of importance 1 to 5.

Rank	REASONS FOR LIVING	Rank	REASONS FOR DYING

I wish to live to the following extent: Not at all: 0 1 2 3 4 5 6 7 8 : Very much

I wish to die to the following extent: Not at all: 0 1 2 3 4 5 6 7 8 : Very much

The one thing that would help me no longer feel suicidal would be:

CAMS Suicide Status Form (SSF-IV-R) Initial Session

Section B (Clinician):

Y N Suicide ideation Describe: _____

☐ Frequency _____ per day _____ per week _____ per month
☐ Duration _____ seconds _____ minutes _____ hours

Y N Suicide plan When: _____

Where: _____

How: _____ Access to means Y N

How: _____ Access to means Y N

Y N Suicide preparation Describe: _____

Y N Suicide rehearsal Describe: _____

Y N History of suicidal behaviors

- Single attempt Describe: _____
- Multiple attempts Describe: _____

Y N Impulsivity Describe: _____

Y N Substance abuse Describe: _____

Y N Significant loss Describe: _____

Y N Relationship problems Describe: _____

Y N Burden to others Describe: _____

Y N Health/pain problems Describe: _____

Y N Sleep problems Describe: _____

Y N Legal/financial issues Describe: _____

Y N Shame Describe: _____

Section C (Clinician):

TREATMENT PLAN (Refer to Sections A & B)

Problem #	Problem Description	Goals and Objectives	Interventions	Duration
1	<i>Self-Harm Potential</i>	<i>Safety and Stability</i>	<i>Stabilization</i> <i>Plan Completed</i> ☐	
2				
3				

YES _____ NO _____ Patient understands and concurs with treatment plan?

YES _____ NO _____ Patient at imminent danger of suicide (hospitalization indicated)?

Patient Signature

Date

Clinician Signature

Date

Patient: _____ Clinician: _____ Date: _____ Time: _____

Rate each item according to how you feel right now.

- Low pain: 1 2 3 4 5 :High pain

- Low stress:** 1 2 3 4 5 **:High stress**

- Low agitation: 1 2 3 4 5 :High agitation**

- Low hopelessness: 1 2 3 4 5 :High hopelessness**

- Low self-hate: 1 2 3 4 5 :High self-hate**

- In the past week: Suicidal Thoughts/Feelings Y__ N__ Managed Thoughts/Feelings Y__ N__ Suicidal Behavior Y__ N__**

****Complete SSF Outcome Form at 3rd consecutive resolution session****

Patient Status:

☐ Discontinued treatment ☐ No show ☐ Cancelled ☐ Hospitalization ☐ Referred/Other: _____

Problem #	Problem Description	Goals and Objectives	Interventions	Duration
1	Self-Harm Potential	Safety and Stability	Stabilization Plan Updated ☐	
2				
3				

Patient Signature

Date _____

Clinician Signature

Date _____

Section C (Clinician Post-Session Evaluation):

MENTAL STATUS EXAM (circle appropriate items):

ALERTNESS: ALERT DROWSY LETHARGIC STUPOROUS
 OTHER: _____
ORIENTED TO: PERSON PLACE TIME REASON FOR EVALUATION
MOOD: EUTHYMIC ELEVATED DYSPHORIC AGITATED ANGRY
AFFECT: FLAT BLUNTED CONSTRICTED APPROPRIATE LABILE
THOUGHT CONTINUITY: CLEAR & COHERENT GOAL-DIRECTED TANGENTIAL CIRCUMSTANTIAL
 OTHER: _____
THOUGHT CONTENT: WNL OBSESSIONS DELUSIONS IDEAS OF REFERENCE BIZARRENESS MORBIDITY
 OTHER: _____
ABSTRACTION: WNL NOTABLY CONCRETE
 OTHER: _____
SPEECH: WNL RAPID SLOW SLURRED IMPOVERISHED INCOHERENT
 OTHER: _____
MEMORY: GROSSLY INTACT
 OTHER: _____
REALITY TESTING: WNL
 OTHER: _____

NOTABLE BEHAVIORAL OBSERVATIONS: _____

DIAGNOSTIC IMPRESSIONS/DIAGNOSIS (DSM/ICD DIAGNOSES):

PATIENT'S OVERALL SUICIDE RISK LEVEL (check one and explain):

- ☐ **MILD (WTL/RFL)** **Explanation:** _____
☐ **MODERATE (AMB)** _____
☐ **HIGH (WTD/RFD)** _____

CASE NOTES:

Next Appointment Scheduled: _____ Treatment Modality: _____

Clinician Signature

Date

CAMS Suicide Status Form (SSF-IV-R) Outcome/Disposition Final Session

Patient: _____ Clinician: _____ Date: _____ Time: _____

Section A (Patient):

Rate each item according to how you feel right now.

1) RATE PSYCHOLOGICAL PAIN (*hurt, anguish, or misery in your mind, not stress, not physical pain*):

Low pain: 1 2 3 4 5 :High pain

2) RATE STRESS (*your general feeling of being pressured or overwhelmed*):

Low stress: 1 2 3 4 5 :High stress

3) RATE AGITATION (*emotional urgency; feeling that you need to take action; not irritation; not annoyance*):

Low agitation: 1 2 3 4 5 :High agitation

4) RATE HOPELESSNESS (*your expectation that things will not get better no matter what you do*):

Low hopelessness: 1 2 3 4 5 :High hopelessness

5) RATE SELF-HATE (*your general feeling of disliking yourself; having no self-esteem; having no self-respect*):

Low self-hate: 1 2 3 4 5 :High self-hate

6) RATE OVERALL RISK OF
SUICIDE:

Extremely low risk: 1 2 3 4 5 :Extremely high risk
(will not kill self) (will kill self)

In the past week: Suicidal Thoughts/Feelings Y__ N__ Managed Thoughts/Feelings Y__ N__ Suicidal Behavior Y__ N__

Were there any aspects of your treatment that were particularly helpful to you? If so, please describe these. Be as specific as possible.

What have you learned from your clinical care that could help you if you became suicidal in the future?

Section B (Clinician):

Third consecutive session of resolved suicidality: ____ Yes ____ No (if no, continue CAMS tracking)

****Resolution of suicidality, if for third consecutive week: current overall risk of suicide <3; in past week: no suicidal behavior and effectively managed suicidal thoughts/feelings**

OUTCOME/DISPOSITION (Check all that apply):

____ Continuing outpatient psychotherapy ____ Inpatient hospitalization

____ Mutual termination ____ Patient chooses to discontinue treatment (unilaterally)

____ Referral to: _____

____ Other. Describe: _____

Next Appointment Scheduled (if applicable): _____

Patient Signature

Date

Clinician Signature

Date

Section C (Clinician Outcome Evaluation):

MENTAL STATUS EXAM (circle appropriate items):

ALERTNESS: ALERT DROWSY LETHARGIC STUPOROUS
 OTHER: _____
ORIENTED TO: PERSON PLACE TIME REASON FOR EVALUATION
MOOD: EUTHYMIC ELEVATED DYSPHORIC AGITATED ANGRY
AFFECT: FLAT BLUNTED CONSTRICTED APPROPRIATE LABILE
THOUGHT CONTINUITY: CLEAR & COHERENT GOAL-DIRECTED TANGENTIAL CIRCUMSTANTIAL
 OTHER: _____
THOUGHT CONTENT: WNL OBSESSIONS DELUSIONS IDEAS OF REFERENCE BIZARRENESS MORBIDITY
 OTHER: _____
ABSTRACTION: WNL NOTABLY CONCRETE
 OTHER: _____
SPEECH: WNL RAPID SLOW SLURRED IMPOVERISHED INCOHERENT
 OTHER: _____
MEMORY: GROSSLY INTACT
 OTHER: _____
REALITY TESTING: WNL
 OTHER: _____

NOTABLE BEHAVIORAL OBSERVATIONS: _____

DIAGNOSTIC IMPRESSIONS/DIAGNOSIS (DSM/ICD DIAGNOSES):

PATIENT'S OVERALL SUICIDE RISK LEVEL (check one and explain):

- ☐ **LOW (WTL/RFL)** **Explanation:** _____
☐ **MODERATE (AMB)** _____
☐ **HIGH (WTD/RFD)** _____

CASE NOTES:

Clinician Signature

Date

CAMS Suicide Status Form (SSF-IV-R) STABILIZATION PLAN

Ways to reduce access to lethal means:

1. _____
2. _____
3. _____

Things I can do to cope differently when I am in a suicide crisis (consider crisis card):

1. _____
2. _____
3. _____
4. _____
5. _____

6. Life or death emergency contact number: _____

People I can call for help or to decrease my isolation:

1. _____
2. _____
3. _____

Attending treatment as scheduled:

Potential Barrier:

Solutions I will try:

1. _____
2. _____

Section D (Clinician Post-Session Evaluation):

MENTAL STATUS EXAM (circle appropriate items):

ALERTNESS: ALERT DROWSY LETHARGIC STUPOROUS
OTHER: _____
ORIENTED TO: PERSON PLACE TIME REASON FOR EVALUATION
MOOD: EUTHYMIC ELEVATED DYSPHORIC AGITATED ANGRY
AFFECT: FLAT BLUNTED CONSTRICTED APPROPRIATE LABILE
THOUGHT CONTINUITY: CLEAR & COHERENT GOAL-DIRECTED TANGENTIAL CIRCUMSTANTIAL
OTHER: _____
THOUGHT CONTENT: WNL OBSESSIONS DELUSIONS IDEAS OF REFERENCE BIZARRENESS MORBIDITY
OTHER: _____
ABSTRACTION: WNL NOTABLY CONCRETE
OTHER: _____
SPEECH: WNL RAPID SLOW SLURRED IMPOVERISHED INCOHERENT
OTHER: _____
MEMORY: GROSSLY INTACT
OTHER: _____
REALITY TESTING: WNL
OTHER: _____

NOTABLE BEHAVIORAL OBSERVATIONS: _____

DIAGNOSTIC IMPRESSIONS/DIAGNOSIS (DSM/ICD DIAGNOSES):

PATIENT'S OVERALL SUICIDE RISK LEVEL (check one and explain):

- Will to Live - Reasons*
☐ **LOW (WTL/RFL)** **Explanation:**
are they ambivalent
☐ **MODERATE (AMB)** _____

☐ **HIGH (WTD/RFD)** _____

CASE NOTES:

Next Appointment Scheduled: _____ Treatment Modality: _____

Clinician Signature

Date